

A note on continuous ensemble expansions of quantum states

Romàn R. Zapatrin

Friedmann Lab. for Theoretical Physics, SPb EF University,

Griboyedova 30–32, 191023, St.Petersburg, Russia;

e-mail: zapatrin@rusmuseum.ru

Abstract

Generalizing the notion of relative entropy, the difference between *a priori* and *a posteriori* relative entropy for quantum systems is drawn. The former, known as quantum relative entropy, is associated with quantum states recognition. The latter—*a posteriori* relative quantum entropy is introduced and shown to be related with state reconstruction due to the following property: given a density operator ρ , ensembles of pure states with Gibbs distribution with respect to the defined distance are proved to represent the initial state ρ up to an amount of white noise (completely mixed state) which can be made arbitrary small.

In classical probability the RELATIVE ENTROPY (or Kullback-Leibler distance) $S(\rho||\sigma)$ of a distribution $\rho = \{p_1, \dots, p_n\}$ with respect to another distribution $\sigma = \{q_1, \dots, q_n\}$ is defined as

$$S(\rho||\sigma) = \begin{cases} -\sum_k p_k \log\left(\frac{p_k}{q_k}\right) & \text{if } \text{supp } \rho \subseteq \text{supp } \sigma \\ +\infty & \text{otherwise} \end{cases} \quad (1)$$

Usually, this notion is generalized in quantum information theory (see, *e.g.* [5] for a review) by analogy with von Neumann entropy, namely, the sum is replaced by operator trace. For two density operators ρ and σ the quantum relative entropy $S(\rho||\sigma)$ reads:

$$S(\rho||\sigma) = \begin{cases} -\text{Tr}[\rho(\log \rho - \log \sigma)] & \text{if } \text{supp } \rho \subseteq \text{supp } \sigma \\ +\infty & \text{otherwise} \end{cases} \quad (2)$$

Relative entropy and state recognition. Although relative entropy does not satisfy the triangle inequality and is therefore not a ‘true’ metric, it is a nonnegative convex function of $\{p_k\}$ and equals zero only if the distributions are equal, $\sigma = \rho$. Its relevance to distinguishing probability distributions is vindicated by the Sanov’s theorem [1] which states that the probability $P_N(\rho|\sigma)$ that the state σ passes the test determining if the state is ρ tends to

$$P_N(\rho|\sigma) \rightarrow \text{const} \cdot e^{-NS(\rho||\sigma)} \quad (3)$$

as N (the number of checked samples of σ) tends to ∞ . A quantum analog of Sanov theorem was also proved [2]. This shows that both classical and quantum relative entropy is an adequate tool to *recognize* states.

Relative entropy from the operationalistic perspective. In this note I give a quantum *operationalistic* analog of Kullback-Leibler distance (1). For that, note that each probability q_k can be treated in two ways

- *a priori*, that is as the probability of the event associated with the value q_k of σ to occur
- *a posteriori*, that is, as the probability of the event associated with the value p_k of ρ to occur provided the system was in state σ

Coming to quantum formula according to *a priori* interpretation, we get the standard quantum relative entropy (2) of ρ with respect to σ . Following the second analogy, we get the relative entropy of ρ with respect to the post-measurement state, which is called LÜDERS STATE [4], it reads

$$\rho_L[\sigma] = \sum_{k=1}^n |\mathbf{e}_k\rangle\langle\mathbf{e}_k| \sigma |\mathbf{e}_k\rangle\langle\mathbf{e}_k| \quad (4)$$

and describes the state of the system which was initially in state σ after the measurement associated with the density operator ρ was carried out. Now we can introduce the distance between the states—the *a posteriori* quantum generalization of relative entropy—as follows

$$L(\rho||\phi) = S(\rho||\rho_L[\sigma]) \quad (5)$$

Note that the value of this distance may be finite when $\sigma = |\phi\rangle\langle\phi|$ is a pure state and ρ is mixed, which is never the case for quantum relative entropy. For further purposes the explicit expression—an analog of (2)—for the distance function (5) between a pure state ϕ and mixed state $\rho = \sum_{k=1}^n p_k |\mathbf{e}_k\rangle\langle\mathbf{e}_k|$ is to be written down:

$$L(\rho||\phi) = \begin{cases} -\sum_k p_k \log \frac{p_k}{|\langle\mathbf{e}_k|\phi\rangle|^2} & \text{if } \phi \text{ has nonzero overlap with } \rho \\ +\infty & \text{otherwise} \end{cases} \quad (6)$$

—a vector ϕ is said to have [3] a NONZERO OVERLAP with ρ if $\langle\phi|\rho|\phi\rangle \neq 0$.

Reconstructing quantum states by Gibbs ensembles. Given a full-range density operator ρ in $\mathbf{H} = \mathbb{C}^n$, form a kind of Gibbs ensemble of pure states treating $L(\rho||\phi)$ as distance:

$$\mu_\rho(\phi) = K \exp\left(-\frac{n}{\varepsilon} L(\rho||\phi)\right) \quad (7)$$

where $\varepsilon > 0$ is a small parameter and K is a normalization constant with respect to Haar measure dS on the set $\mathbb{C}S^n$ of unit vectors in $\mathbf{H}$. This ensemble is associated with a density operator

$$I(\rho) = \int_{\mathbb{C}S^n} \mu_\rho(\phi) |\phi\rangle\langle\phi| dS \quad (8)$$

and the following RECONSTRUCTION FORMULA holds:

$$\int_{\mathbb{C}S^n} \mu_\rho(\phi) |\phi\rangle\langle\phi| dS = (1 - \varepsilon)\rho + \varepsilon\Lambda \quad (9)$$

where $\Lambda = \frac{1}{n}$ is the density matrix of white noise—a completely mixed state. The proof of this formula is a routine integration, see Appendix for details and the explicit expression for the normalization constant.

Limit distribution. The limit distribution in (9) corresponds to an ensemble $\mathcal{E}$ whose density matrix is exactly ρ , let us describe it explicitly. According to (9), the support of the limit distribution is the set of unit vectors for which the *a posteriori* distance from ρ is zero, that is, $\text{supp}(\mathcal{E}) = \{\phi_{\theta_1, \dots, \theta_n} \mid L(\rho \parallel \phi_{\theta_1, \dots, \theta_n}) = 0\}$ with

$$\phi_{\theta_1, \dots, \theta_n} = \sum_{k=1}^n e^{-\theta_k} \sqrt{p_k} |\mathbf{e}_k\rangle$$


where $\theta_1, \dots, \theta_n = 0 \dots 2\pi$. Therefore the resulting distribution is uniform over the n -torus $S^1 \times \dots \times S^1$ and, as a consequence, any full-range density operator ρ can be represented as the following UNIFORM ENSEMBLE

$$\rho = \frac{1}{(2\pi)^n} \int_0^{2\pi} \dots \int_0^{2\pi} |\phi_{\theta_1, \dots, \theta_n}\rangle \langle \phi_{\theta_1, \dots, \theta_n}| d\theta_1 \dots d\theta_n \quad (10)$$


An example. Consider a density operator ρ whose spectral decomposition is

$$\rho = \frac{1}{4} |0\rangle \langle 0| + \frac{3}{4} |1\rangle \langle 1| \quad (11)$$


in two-dimensional real space $\mathbf{H} = \mathbb{R}^2$. The picture below shows three expansions of ρ (note that in case of real state space the uniform ensemble (10) is discrete as the phase multiples are ± 1)



Gibbs ensemble (9)



Uniform ensemble (10)



Classical ensemble (11)

Acknowledgments. The attention to work offered by the participants of the Friedmann seminar on theoretical physics (St.Petersburg), in particular, to S.Krasnikov and A.Lobashev is highly appreciated.

References

- [1] Cover, T. M. and Thomas, J. A., *Elements of Information Theory*. New York: Wiley, 1991
- [2] F. Hiai, D. Petz, The proper formula for relative entropy and its asymptotics in quantum probability. *Communications in Mathematical Physics*, **143** 99 (1991)
- [3] M.Horodecki, Aditi Sen De, Ujjwal Sen, Quantification of quantum correlation of ensemble of states, quant-ph/0310100

- [4] G.Lüders, Über die Zustandsänderung durch den Meßprozeß, *Ann. Physik*, **8** (6) 322 (1951); English translation ‘Regarding the state-change due to the measurement process’ by K.A.Kirkpatrick is available at [quant-ph/0403007](#)
- [5] M.Ohya, D.Petz, *Quantum entropy and its use*, Springer, 1993

Appendix: the proof of formula (9)

Introduce the following numerical integral

$$I_n^{a_1 \dots a_n} = \int_{z \in \mathbb{C}S_n} \prod_{k=1}^n |\langle \mathbf{e}_k | z \rangle|^{2a_k} dS \quad (12)$$

for which the following lemma holds

Lemma 1. For $a_1, \dots, a_n \geq 0$ and $\langle \mathbf{e}_k | \mathbf{e}_j \rangle = \delta_{kj}$

$$I_n^{a_1 \dots a_n} = \frac{2\pi^n \prod_{k=1}^n \Gamma(a_k + 1)}{\Gamma(n + \sum_{k=1}^n a_k)} \quad (13)$$

Proof. Represent $\mathbb{C}^n$ as real space $\mathbb{R}^{2n}$ with coordinates $(r_1, \phi_1; \dots; r_n, \phi_n)$ so that $\langle \mathbf{e}_k | z \rangle = z_k = r_k e^{i\phi_k}$. Write down the integral (12) as

$$I_n^{a_1 \dots a_n} = (2\pi)^n \int_{r_1^2 + \dots + r_n^2 = 1} \prod_{k=1}^n r_k^{2a_k} dS$$

Getting rid of $r_1 = \sqrt{1 - r_2^2 - \dots - r_n^2}$ we obtain

$$I_n^{a_1 \dots a_n} = (2\pi)^n \cdot \int_{r_2^2 + \dots + r_n^2 \leq 1} \left(1 - \sum_{j=2}^n r_j^2\right)^{a_1} \left(\prod_{k=2}^n r_k^{2a_k+1} dr_k\right)$$

Now, integrating over r_n , we obtain

$$I_n^{a_1 \dots a_n} = (2\pi)^n \cdot \int_0^1 dr_n r_n^{2a_n+1} \int_{r_2^2 + \dots + r_{n-1}^2 \leq 1 - r_n^2} \left(1 - r_n^2 - \sum_{j=2}^{n-1} r_j^2\right)^{a_1} \left(\prod_{k=2}^{n-1} r_k^{2a_k+1} dr_k\right)$$

which, under the substitution $\xi_k = r_k / \sqrt{1 - r_n^2}$ for $k = 2, \dots, n-1$ and $r_n = \sin \alpha$ reads

$$\begin{aligned} I_n^{a_1 \dots a_n} &= (2\pi)^n \cdot \int_0^{\pi/2} d\sin \alpha \sin^{2a_n+1} \alpha \cos^{2a_1} \alpha (\cos \alpha)^{2(\sum_{k=2}^{n-1} a_k) + 2(n-2)} \times \\ &\quad \times \int_{\xi_2^2 + \dots + \xi_{n-1}^2 \leq 1} \left(1 - \sum_{j=2}^{n-1} \xi_j^2\right)^{a_1} \left(\prod_{k=2}^{n-1} \xi_k^{2a_k+1} d\xi_k\right) = \\ &= 2\pi \int_0^{\pi/2} \sin^{2a_n+1} (\cos \alpha)^{2(\sum_{k=1}^{n-1} a_k + n-2) + 1} d\alpha \cdot I_{n-1}^{a_1 \dots a_{n-1}} \end{aligned}$$

Using L egendre's formula

$$\int_0^{\pi/2} \sin^{2\alpha+1} x \cos^{2\beta+1} x dx = \frac{1}{2} \cdot \frac{\Gamma(\alpha+1)\Gamma(\beta+1)}{\Gamma(\alpha+\beta+2)} \quad (14)$$

we obtain the following recurrent expression

$$I_n^{a_1 \dots a_n} = \pi \cdot \frac{\Gamma(a_n+1)\Gamma(n-1+\sum_{k=1}^{n-1} a_k)}{\Gamma(n+\sum_{k=1}^n a_k)} \cdot I_{n-1}^{a_1 \dots a_{n-1}} \quad (15)$$

Direct calculations show that the formula (13) satisfies the recurrent expression (15), and it remains to prove the induction base. Do it for $n = 1$:

$$I_1^p = \int_0^{2\pi} d\phi \cdot 1^p = 2\pi$$

which also accords with (13). This completes the proof. $\square$

Now let $A = \sum_{k=1}^n a_k |\mathbf{e}_k\rangle\langle\mathbf{e}_k|$ be a positive self-adjoint operator in $\mathbf{H} = \mathbb{C}^n$. Introduce the operator integral

$$I(A) = \int_{\mathbb{C}S^n} \left(\prod_{k=1}^n |\langle \mathbf{e}_k | \phi \rangle|^{2a_k} \right) |\phi\rangle\langle\phi| dS \quad (16)$$

for which the following lemma holds.

Lemma 2. *For any positive self-adjoint operator A in $\mathbf{H}$*

$$I(A) = \frac{2\pi^n \prod_{k=1}^n \Gamma(1 + a_k)}{\Gamma(1 + n + \text{Tr } A)} (A + \mathbb{I}) \quad (17)$$

Proof. First prove that $I(A)$ is diagonal in the eigenbasis of A . For any $j \neq k$ the appropriate matrix element reads

$$\langle \mathbf{e}_j | I(A) | \mathbf{e}_k \rangle = (2\pi)^{n-2} \iint_0^{2\pi} e^{i(\phi_k - \phi_j)} d\phi_j d\phi_k \int_{r_1^2 + \dots + r_n^2 = 1} \prod_i r^{2a_i} dS = 0$$

Now calculate the diagonal elements

$$\langle \mathbf{e}_k | I(A) | \mathbf{e}_k \rangle = \int_{\mathbb{C}S_n} \left(\prod_{i=1}^n |\langle \mathbf{e}_i | \phi \rangle|^{2a_i} \right) |\langle \mathbf{e}_k | \phi \rangle|^2 dS = I_n^{a_1 \dots (a_k+1) \dots a_n}$$

In the meantime it follows from (13) that

$$I_n^{a_1 \dots (a_k+1) \dots a_n} = I_n^{a_1 \dots a_n} \frac{a_k + 1}{\Gamma(n + 1 + \sum_i a_i)}$$

then $I(A) = \sum_k \langle \mathbf{e}_k | I(A) | \mathbf{e}_k \rangle \cdot |\mathbf{e}_k\rangle\langle\mathbf{e}_k|$, hence

$$I(A) = \frac{2\pi^n \prod \Gamma(a_i + 1)}{\Gamma(n + 1 + \sum a_i)} \sum_{k=1}^n (a_k + 1) |\mathbf{e}_k\rangle\langle\mathbf{e}_k|$$

which completes the proof. $\square$

Denote by Λ the density matrix of the white noise—a completely mixed state

$$\Lambda = \frac{1}{n} \mathbb{I} = \frac{1}{n} \sum_{k=1}^n |\mathbf{e}_k\rangle\langle\mathbf{e}_k|$$

then the exact version of formula (9) is stated by the following theorem.

Theorem 3. Let $\rho = \sum_{k=1}^n p_k |\mathbf{e}_k\rangle\langle\mathbf{e}_k|$ be a positive ($p_k > 0$) operator in $\mathbf{H} = \mathbb{C}^n$. Then for any $\varepsilon \in (0, 1)$

$$(1 - \varepsilon)\rho + \varepsilon\Lambda = \frac{\Gamma\left(1 + \frac{n}{\varepsilon}\right) e^{-\frac{n(1-\varepsilon)}{\varepsilon}S(\rho)}}{2\pi^n \prod_{k=1}^n \Gamma\left(1 + \frac{n(1-\varepsilon)p_k}{\varepsilon}\right)} \int_{\mathbb{C}S^n} e^{-\frac{n(1-\varepsilon)}{\varepsilon}L(\rho||\phi)} |\phi\rangle\langle\phi| dS \quad (18)$$

Proof. Consider the formula (17) and let $A = \frac{n(1-\varepsilon)}{\varepsilon}\rho$, then $a_k = \frac{n(1-\varepsilon)p_k}{\varepsilon}$ and

$$\frac{2\pi^n \prod_{k=1}^n \Gamma(1 + a_k)}{\Gamma(1 + n + \text{Tr } A)} \left(\frac{n(1-\varepsilon)}{\varepsilon}\rho + \mathbb{I} \right) = \int_{\mathbb{C}S^n} \left(\prod_{k=1}^n |\langle \mathbf{e}_k | \phi \rangle|^2 \frac{n(1-\varepsilon)p_k}{\varepsilon} \right) |\phi\rangle\langle\phi| dS$$

thus, taking into account that $\text{Tr } A = \frac{n(1-\varepsilon)}{\varepsilon}$, we have

$$\frac{n(1-\varepsilon)}{\varepsilon}\rho + \mathbb{I} = \frac{\Gamma\left(1 + \frac{n}{\varepsilon}\right)}{2\pi^n \prod_{k=1}^n \Gamma\left(\frac{n(1-\varepsilon)p_k}{\varepsilon} + 1\right)} \int_{\mathbb{C}S^n} e^{\frac{n(1-\varepsilon)}{\varepsilon} \sum_{k=1}^n p_k \log |\langle \mathbf{e}_k | \phi \rangle|^2} |\phi\rangle\langle\phi| dS$$

Using the definition (6) we obtain $\frac{n(1-\varepsilon)}{\varepsilon}\rho + \mathbb{I} =$

$$= \frac{\Gamma\left(1 + \frac{n}{\varepsilon}\right) e^{\frac{n(1-\varepsilon)}{\varepsilon} \sum_{k=1}^n p_k \log p_k}}{2\pi^n \prod_{k=1}^n \Gamma\left(1 + \frac{n(1-\varepsilon)p_k}{\varepsilon}\right)} \int_{\mathbb{C}S^n} e^{-\frac{n(1-\varepsilon)}{\varepsilon} \sum_{k=1}^n p_k (\log p_k - \log |\langle \mathbf{e}_k | \phi \rangle|^2)} |\phi\rangle\langle\phi| dS$$

from which the formula (18) follows directly. $\square$

I emphasize that the formula (18) is exact. For small ε we assume $1 - \varepsilon \simeq 1$ and obtain the formula (9).